

2011-2012 Missouri School Year Immunization Requirements

Revised 03/15/11

GR	DTaP/DTP/DT/Td/Tdap	POLIO	MEASLES	MUMPS	RUBELLA	HEPATITIS B	VARICELLA
	4+ DOSES	3+ DOSES	2 DOSES	2 DOSES	2 DOSES	3+ DOSES	2 DOSES
K-1	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses.	Last dose on or after 4th birthday and at least 6 months after previous dose. If 4 doses before 4 years of age, a 5th dose should be given at 4 to 6 years of age.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Last dose must be administered on or after 168 days of age. (There is a 4 day grace period which would make it 164 days.)	Dose 1 on or after first birthday and a minimum of 28 days between dose 1 and dose 2. (Recommended 3 months) or a statement signed by MD or DO indicating the month and year the child had chickenpox disease.
	4+ DOSES	3+ DOSES	2 DOSES	2 DOSES	2 DOSES	3+ DOSES *	1 DOSE
2-6	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses.	Last dose on or after fourth (4th) birthday. If a combination of IPV/OPV is received, four (4) doses are required. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Last dose must be administered on or after 168 days of age. (There is a 4 day grace period which would make it 164 days.)	On or after first birthday or a statement signed by parent/guardian or physician indicating the month and year the child had chickenpox disease.
	4+ DOSES	3+ DOSES	2 DOSES	2 DOSE	2 DOSE	3+ DOSES *	
7	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses.	Last dose on or after fourth (4th) birthday. If a combination of IPV/OPV is received, four (4) doses are required. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Last dose must be administered on or after 168 days of age. (There is a 4 day grace period which would make it 164 days.)	
	4+ DOSES	3+ DOSES	2 DOSES	2 DOSES	2 DOSES	3+ DOSES	
8	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses. 1 Tdap required unless had a Td Booster within the last two years.	Last dose on or after fourth (4th) birthday. If a combination of IPV/OPV is received, four (4) doses are required. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Dose 1 - Initial visit. Dose 2 - 28 days after Dose 1. Dose 3 - Minimum 56 days between dose 2 & 3. Cannot complete series in less than 4 months. There is a 4 day grace period.	
	4 DOSES	3+ DOSES	2 DOSES	1 DOSE	1 DOSE	3+ DOSES	
9	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses. 1 Tdap required unless had a Td Booster within the last two years.	Last dose on or after fourth (4th) birthday. If a combination of IPV/OPV is received, four (4) doses are required. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Dose 1 - Initial visit. Dose 2 - 28 days after Dose 1. Dose 3 - Minimum 56 days between dose 2 & 3. Cannot complete series in less than 4 months. There is a 4 day grace period.	** There is a 4 day grace period for all types of vaccines
	4 DOSES	3+ DOSES	2 DOSES	1 DOSE	1 DOSE	3+ DOSES	
10-12	Last dose on or after fourth (4th) birthday and last dose of pertussis before seventh (7th) birthday. Maximum needed: six (6) doses. Td Booster or Tdap required 10 year after last does of DTaP, DTP, DT, Td.	Last dose on or after fourth (4th) birthday. If a combination of IPV/OPV is received, four (4) doses are required. Maximum needed: four (4) doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	On or after first (1st) birthday. Twenty-eight (28) days between the two doses.	Dose 1 - Initial visit. Dose 2 - 28 days after Dose 1. Dose 3 - Minimum 56 days between dose 2 & 3. Cannot complete series in less than 4 months. There is a 4 day grace period.	*Hep B note= Dosage intervals between dose1,2,3 are the same for all grades k-12